

THE DISCRIMINANT OF A SPACE CURVE IS STABLE

SEAN TIMOTHY PAUL

ABSTRACT. We prove that the discriminant of a nonsingular space curve of genus $g \geq 2$ is stable with respect to the standard action of the special linear group.

Historically the construction of the moduli space of curves of genus g was carried out by David Mumford in his epoch making treatise *Geometric Invariant Theory*. The basic first step in the construction was to verify the stability of either the Cayley-Chow form or the Hilbert point of a (smooth) complex projective-algebraic curve.

Theorem. (Mumford [MFK94]) Let $X \rightarrow \mathbb{P}^N$ be a nonsingular space curve of genus $g \geq 2$ embedded by a very ample complete linear system. Assume that $d = \deg(X) \geq 2g$. Let R_X denote the **Cayley-Chow form** of X . Then R_X is stable with respect to the action of $G = SL(N+1, \mathbb{C})$.

Theorem. (Geiseker [Gie82]) Let $X \rightarrow \mathbb{P}^N$ be a nonsingular space curve of genus $g \geq 2$ embedded by a very ample complete linear system. Assume that $d = \deg(X) \geq 2g$. Let $\mathcal{H}_m(X)$ denote the m th **Hilbert point** of X . Then there exists a uniform constant m_0 such that $\mathcal{H}_m(X)$ is stable with respect to the action of G for all $m \geq m_0$.

The purpose of this note is to establish the corresponding statement for the dual of a nonsingular algebraic curve.

Theorem 0.1. Let $X \subset \mathbb{P}^N$ be a nonsingular space curve of genus $g \geq 2$ embedded by a very ample complete linear system. Assume that $d = \deg(X) \geq 2g$. Let Δ_X denote the X -**discriminant** (the defining polynomial of the dual variety $X^\vee$). Then Δ_X is stable with respect to the action of G .

0.1. Resultants and Discriminants. Let $X \rightarrow \mathbb{P}^N$ be a complex algebraic curve. Let R_X denote the Cayley-Chow form of $X \rightarrow \mathbb{P}^N$ and let Δ_X be the X -discriminant (see [GKZ94]). For our purpose we must normalize these polynomials

$$(0.1) \quad R := R_X^{\deg(\Delta_X)}, \quad \Delta := \Delta_X^{\deg(R_X)}.$$

It is well known that up to scaling these lie in finite dimensional (irreducible) G -modules $\mathbb{E}_{\lambda_\bullet}$ and $\mathbb{E}_{\mu_\bullet}$ respectively. The reader can consult [Pau11] for details on these modules.

0.2. The Mabuchi Energy. Let (X, ω) be a closed Kähler manifold of dimension n , where ω denotes the Kähler form. The space of Kähler potentials will be denoted by $\mathcal{H}_\omega$

$$\mathcal{H}_\omega := \left\{ \varphi \in C^\infty(X) \mid \omega_\varphi := \omega + \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \varphi > 0 \right\}.$$

Definition 1. (Mabuchi, [Mab86]) The **K-Energy** of (X, ω) is the map $\nu_\omega : \mathcal{H}_\omega \longrightarrow \mathbb{R}$ given by the following expression

$$\nu_\omega(\varphi) := -\frac{1}{V} \int_0^1 \int_X \dot{\varphi}_t (\text{Scal}(\varphi_t) - \mu) \omega_t^n dt.$$

φ_t is a smooth path in $\mathcal{H}_\omega$ joining 0 with φ , $\text{Scal}(\varphi_t)$ denotes the scalar curvature of the metric $\omega_t := \omega + \sqrt{-1} \partial \bar{\partial} \varphi_t$, μ denotes the average of the scalar curvature of ω , and V is the volume.

It is well known that the K-Energy does not depend on the path chosen (see [Mab86]). Suppose that ω represents a multiple of the *canonical class*

$$(0.2) \quad \text{Ric}(\omega) = \frac{\mu}{n} \omega + \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} h_\omega.$$

In this case there is the following well known direct formula for the K-energy.

$$(0.3) \quad \begin{aligned} \nu_\omega(\varphi) &= \int_X \log \left(\frac{\omega_\varphi^n}{\omega^n} \right) \frac{\omega_\varphi^n}{V} - \frac{\mu}{n} (I_\omega(\varphi) - J_\omega(\varphi)) - \frac{1}{V} \int_X h_\omega(\omega_\varphi^n - \omega^n) \\ J_\omega(\varphi) &:= \frac{1}{V} \int_X \sum_{i=0}^{n-1} \frac{\sqrt{-1}}{2\pi} \frac{i+1}{n+1} \partial \varphi \wedge \bar{\partial} \varphi \wedge \omega^i \wedge \omega_\varphi^{n-i-1} \end{aligned}$$

$$I_\omega(\varphi) := \frac{1}{V} \int_X \varphi (\omega^n - \omega_\varphi^n).$$

When X is a compact Riemann surface of genus g (i.e. an algebraic curve) equipped with some Kähler metric ω the complicated expression above reduces to

$$(0.4) \quad \nu_\omega(\varphi) = \int_X \log \left(\frac{\omega_\varphi}{\omega} \right) \frac{\omega_\varphi}{V} + \frac{(2g-2)}{2V^2} \int_X |\nabla \varphi|^2 \omega - \frac{1}{V} \int_X h_\omega(\omega_\varphi - \omega).$$

Proposition 0.1. Let X be a smooth algebraic curve of genus $g \geq 1$. Let ω be any Kähler metric on X . Then ν_ω is bounded below on $\mathcal{H}_\omega$.

Proof. The first integral is bounded below by $\frac{-1}{e}$, the second by 0, and the third by $-||h_\omega||_{C^0}$. $\square$

Remark 1. The Mabuchi energy is bounded below when $g = 0$, but this fact is much more difficult to prove.

Now suppose that $X \subset \mathbb{P}^N$ is a smooth subvariety (in particular, a projective curve) and that $\omega = \omega_{FS}|_X$, where ω_{FS} denotes the Fubini-Study Kähler form. Given $\sigma \in G$ we have

$$\sigma^* \omega_{FS} = \omega_{FS} + \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \varphi_\sigma > 0.$$

In this way we produce a set theoretic map

$$G \ni \sigma \rightarrow \varphi_\sigma \in \mathcal{H}_\omega.$$

It is easy to see that φ_σ is given by the formula

$$(0.5) \quad \varphi_\sigma = \log \frac{||\sigma z||^2}{||z||^2}.$$

We define $\nu_\omega(\sigma) := \nu_\omega(\varphi_\sigma)$. In this way we may consider the K -energy as a function on G .

Theorem A . (Paul [Pau11]) Let $X^n \hookrightarrow \mathbb{P}^N$ be a smooth, linearly normal, complex projective variety of degree $d \geq 2$. Then there are continuous norms on $\mathbb{E}_{\mu_\bullet}$ and $\mathbb{E}_{\lambda_\bullet}$ such that the Mabuchi energy restricted to $\mathcal{B}$ is given as follows

$$d^2(n+1)\nu_\omega(\sigma) = \log \frac{\|\sigma \cdot \Delta\|^2}{\|\Delta\|^2} - \log \frac{\|\sigma \cdot R\|^2}{\|R\|^2}.$$

Let $\lambda : \mathbb{C}^* \rightarrow G$ be an algebraic one parameter subgroup of G . We shall refer to such maps as *degenerations*.

Definition 2. Let $\mathbb{V}$ be a rational representation of G . Let $v \in \mathbb{V} \setminus \{0\}$. Let λ be any degeneration in G . The **weight** $w_\lambda(v)$ of λ on v is the unique integer such that

$$\lim_{|t| \rightarrow 0} t^{-w_\lambda(v)} \lambda(t)v \text{ exists in } \mathbb{V} \text{ and is not zero.}$$

The Numerical Criterion of Hilbert and Mumford says that a point $v \in \mathbb{V}$ is (semi)stable if and only if $w_\lambda(v)(\leq) < 0$ for all degenerations λ in G .

Proof. (of Theorem 0.1) An immediate application of Theorem A yields the following.

Proposition 0.2. There is an asymptotic expansion

$$\lim_{|t| \rightarrow 0} \nu_\omega(\lambda(t)) = (w_\lambda(\Delta) - w_\lambda(R)) \log |t|^2 + O(1) \quad t \in \mathbb{C}^*.$$

Proposition 0.1 implies that for every degeneration λ the following inequality holds

$$(0.6) \quad w_\lambda(\Delta) \leq w_\lambda(R).$$

When $d \geq 2g$ Mumfords' theorem (and the numerical criterion) says that $w_\lambda(R) < 0$. Therefore by (0.6) we also have that $w_\lambda(\Delta) < 0$ and the proof is complete. $\square$

Despite the classical nature of the subject the following result seems to be new.

Theorem 0.2. Let $X_F \rightarrow \mathbb{P}^2$ be a nonsingular plane algebraic curve with $\deg(F) \geq 2$. Then the dual curve is semistable with respect to the action of $SL(3, \mathbb{C})$.

We have an interesting corollary of Theorem 0.2.

Corollary 0.1. There exist semistable plane curves with cusp singularities.

REFERENCES

- [Gie82] D. Gieseker. *Lectures on moduli of curves*, volume 69 of *Tata Institute of Fundamental Research Lectures on Mathematics and Physics*. Published for the Tata Institute of Fundamental Research, Bombay, 1982.
- [GKZ94] I. M. Gelfand, M. M. Kapranov, and A. V. Zelevinsky. *Discriminants, resultants, and multidimensional determinants*. Mathematics: Theory & Applications. Birkhäuser Boston Inc., Boston, MA, 1994.
- [Mab86] Toshiki Mabuchi. K -energy maps integrating Futaki invariants. *Tohoku Math. J. (2)*, 38, 1986.
- [MFK94] D. Mumford, J. Fogarty, and F. Kirwan. *Geometric invariant theory*, volume 34 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)]*. Springer-Verlag, Berlin, third edition, 1994.
- [Pau11] Sean Timothy Paul. Hyperdiscriminant polytopes, Chow polytopes, and Mabuchi energy asymptotics. *Annals of Math.*, (<http://annals.math.princeton.edu/articles/2654>), 2011.